

①

X = random variable, $\mathbb{E}X$ = expected value of X

$$\mathbb{E}(X - \mathbb{E}X)^2 = \text{variance} \equiv \text{Var}(X)$$

$$M_X(t) = \mathbb{E}(e^{tX}), t \in \mathbb{R}$$

moment generating function

$$\mathbb{E}X^{p'} = p'\text{th moment}$$

$$\|X\|_p = (\mathbb{E}|X|^p)^{\frac{1}{p}}, p \in (0, \infty)$$

$$\|X\|_\infty = \text{ess sup } |X|.$$

Probability space: $(\Omega, \Sigma, \mathbb{P})$

sample space,
the set of all possible outcomes

set of events, an event being
a set of outcomes in the sample space

↙ A probability function, which, assigns
a probability to each event.

②

$$L^p(\Omega, \Sigma, \mathbb{P}) = \{X : \|X\|_p < \infty\}$$

If $p \in [1, \infty]$, $\|X\|_p$ is a norm
and L^p is a Banach space

please look up what a "norm"
and a "Banach space" is. When
you do, you may encounter other
unfamiliar notions, so you just keep
scrambling!

The exponent $p=2$ is special because in this case,
 L^2 is not only a Banach space, but also a Hilbert space

please look it up!

The inner product on L^2 is given by

$$\langle X, Y \rangle_{L^2} = \mathbb{E}XY, \quad \|X\|_{L^2} = \left(\mathbb{E}|X|^2 \right)^{\frac{1}{2}}$$

(3)

$$\sigma(X) = \|X - \mathbb{E}X\|_2 = \sqrt{\text{Var}(X)}$$

"
standard
deviation

$$\text{cov}(X, Y) = \mathbb{E}((X - \mathbb{E}X)(Y - \mathbb{E}Y)) =$$

"
covariance of
X & Y

$$\langle X - \mathbb{E}X, Y - \mathbb{E}Y \rangle_2$$

gives a geometric interpretation
of covariance:

the more $X - \mathbb{E}X$ and
 $Y - \mathbb{E}Y$ are aligned, the
larger their inner product and
covariance.

④

We now discuss some inequalities.

Definition: We say that a function φ is convex if

$$\varphi(\lambda x + (1-\lambda)y) \leq \lambda \varphi(x) + (1-\lambda)\varphi(y) \quad \forall \lambda \in [0,1]$$

and all $x, y \in \text{domain}(\varphi)$.

Jensen's inequality: X random variable and φ convex, then

$$\varphi(\mathbb{E} X) \leq \mathbb{E} \varphi(X)$$

to be proved later this week

Corollary: If $0 \leq p \leq q \leq \infty$, Work it out!!

$$\|X\|_p \leq \|X\|_q$$

Minkowski: $\|X + Y\|_p \leq \|X\|_p + \|Y\|_p$

if $p \in [1, \infty]$.

to be proved later this week

(5)

Cauchy-Schwarz:

$$|\mathbb{E}XY| \leq \|X\|_2 \|Y\|_2$$

Hölder: $p, q \in (1, \infty)$ $\frac{1}{p} + \frac{1}{q} = 1$, then

$$|\mathbb{E}XY| \leq \|X\|_p \|Y\|_q$$

Cumulative distribution function (CDF)

$$F_X(t) = \mathbb{P}\{X \leq t\}, \quad t \in \mathbb{R}$$

We shall often work w/ $\mathbb{P}\{X > t\} = 1 - F_X(t)$

Key identity: X non-negative random variable.

Then

$$\mathbb{E}X = \int_0^{\infty} \mathbb{P}\{X > t\} dt$$

Proof: $\mathbb{E}X = \mathbb{E} \int_0^{\infty} 1_{\{t < X\}}(t) dt$, where

$$1_{\{t < X\}}(t) = \begin{cases} 1, & t < X \\ 0, & \text{otherwise} \end{cases}$$

⑥

Why is that? Because

$$X = \int_0^X 1 \, dt = \int_0^\infty \mathbf{1}_{\{t \leq X\}}(t) \, dt$$

Going back,

$$\mathbb{E}X = \mathbb{E} \int_0^\infty \mathbf{1}_{\{t \leq X\}}(t) \, dt$$

$$= \int_0^\infty \mathbb{E} \mathbf{1}_{\{t \leq X\}}(t) \, dt = \int_0^\infty \mathbb{P}\{t \leq X\} \, dt.$$

Changing the order of integration and expectation requires proof. The key result is the Fubini-Tonelli theorem.

Markov inequality: X non-negative random variable and $t > 0$,

$$\mathbb{P}\{X \geq t\} \leq \frac{\mathbb{E}X}{t}$$

we shall explore this thoroughly using python

7

Proof of Markov:

Fix $t > 0$. Write $x = x \cdot 1_{\{x \geq t\}} + x \cdot 1_{\{x < t\}}$

It follows that

$$\mathbb{E}X = \mathbb{E}X \cdot 1_{\{x \geq t\}} + \mathbb{E}X \cdot 1_{\{x < t\}}$$

$$\geq \mathbb{E} \cancel{X} \cdot 1_{\{x \geq t\}} = t \mathbb{P}\{X \geq t\}$$

+ O_t

corresponding to
 $\mathbb{E}X \cdot 1_{\{x < t\}}$

We conclude that

$$\mathbb{P}\{X \geq t\} \leq \frac{1}{t} \mathbb{E}X, \text{ as claimed.}$$

Corollary: (Chebyshev) Let X be a random variable w/ mean μ and variance σ^2 . Then for any $t > 0$,

$$\mathbb{P}\{|X - \mu| \geq t\} \leq \frac{\sigma^2}{t^2}$$

~
Please don't take my word
for it and deduce Chebyshev from
Markov.

(8)

Limit theorems:

We know that $\text{Var}(X_1 + \dots + X_N) = \text{Var}(X_1) + \dots + \text{Var}(X_N)$
still independent if X_i s are independent (check!)

If X_i s have the same distribution w/ mean μ and variance σ^2 , then

$$\text{Var}\left(\frac{1}{N} \sum_{i=1}^N X_i\right) = \frac{\sigma^2}{N} \xrightarrow{N \rightarrow \infty} 0$$

In other words, the variance of the sample mean $\frac{1}{N} \sum_{i=1}^N X_i$ of the sample $\{X_1, \dots, X_N\}$ shrinks to 0 as $N \rightarrow \infty$.

This suggests that, for large N , the sample mean should concentrate around the expected value μ . The quantitative version of this statement is known as the

Strong Law of Large Numbers:

Let $X_1, X_2, \dots, X_N$ be independent identically distributed random variables w/ mean μ . Consider the sum

$$S_N = X_1 + X_2 + \dots + X_N$$

⑨

Then as $N \rightarrow \infty$, $\frac{S_N}{N} \rightarrow \mu$ almost surely.

Borel Law of Large Numbers:

Consider random variables $X_1, X_2, \dots, X_n$, independent, identically distributed and assume the values 0 or 1 w/ probability $\frac{1}{2}$.

Let $S_n = \sum_{k=1}^n X_k$. Then $\frac{S_n}{n} \xrightarrow{n \rightarrow \infty} \frac{1}{2}$ w/ probability 1

Hardy and Littlewood improved this result to

$$\limsup_{n \rightarrow \infty} \frac{\left| \frac{S_n}{n} - \frac{1}{2} \right|}{\sqrt{n \log(n)}} < \frac{1}{\sqrt{2}} \quad (1914)$$

and Khinchin improved this to

$$\text{Prob} \left[\limsup_{n \rightarrow \infty} \frac{\left| \frac{S_n}{n} - \frac{1}{2} \right|}{\sqrt{n \log \log n}} < 1 \right] = 1$$

It might be fun to explore these numerically!

(10)

Another absolutely fundamental limit result is the ~~Ande~~berg-Levy Central Limit Theorem:

Consider the standard normal distribution, denoted by $N(0,1)$, w/ density

$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}, \quad x \in \mathbb{R}$$

Let $X_1, X_2, \dots, X_N, \dots$ be a sequence of independent identically distributed random variables. Let

$S_N = X_1 + \dots + X_N$ and normalize it, to obtain a random variable w/ zero mean and unit variance:

$$Z_N = \frac{S_N - \mathbb{E}S_N}{\sqrt{\text{Var}(S_N)}} = \frac{1}{\sigma\sqrt{N}} \sum_{i=1}^N (X_i - \mu)$$

Then, as $N \rightarrow \infty$,

$Z_N \rightarrow N(0,1)$ in distribution.
i.e. convergence of CDFs.

11

Interesting special case: $X_1, X_2, \dots, X_N, \dots$
independent, identically distributed random
variables, taking on values 0 & 1 w/
probability $(1-p)$ & p respectively
(Bernoulli random variables)

Observe that $EX_i = p$ and $\text{Var}(X_i) = p(1-p)$

~ please make these
calculations!

The Central Limit Theorem implies that

if $S_N = X_1 + \dots + X_N$, as before, then

$$\frac{S_N - Np}{\sqrt{Np(1-p)}} \rightarrow N(0, 1) \text{ in distribution}$$

De Moivre-Laplace theorem
(18th century!!)

(12)

Life is not always this "easy." Suppose that X_i takes on 1 w/ probability p_i w/

$\left. \begin{array}{l} p_i \rightarrow 0 \\ N \rightarrow \infty \end{array} \right\}$ so fast that

S_N has mean $O(1)$ instead of $\sim N$.
 $\text{Ber}(p)$

Then the Central Limit Theorem fails! But what is true?

Definition: Z has a poisson distribution w/ parameter λ
 $\left. \begin{array}{l} \text{random variable} \end{array} \right\}$

$Z \sim \text{Pois}(\lambda)$, if it takes values in $\{0, 1, 2, \dots\}$ w/ probability

$$IP\{Z = k\} = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

Theorem: $X_{N,i}$ $1 \leq i \leq N$ independent random variables $X_{N,i} \sim \text{Ber}(p_{N,i})$ and let

$S_N = \sum_{i=1}^N X_{N,i}$. Assume that as $N \rightarrow \infty$,
 $\max_{i \leq N} p_{N,i} \rightarrow 0$ and $ES_N = \sum_{i=1}^N p_{N,i} \rightarrow \lambda < \infty$.

(13)

Then as $N \rightarrow \infty$,

$S_N \rightarrow \text{Pois}(\lambda)$ in distribution.